

PHOSCARE®



Product Information

GENERAL INFORMATION

PHOSCARE is a product produced using new technologies which are different from the traditional phosphorous fertilizer technologies. Phosphorous acid instead of phosphoric acid is used as the phosphorous source. The product derived from using Phosphorous acid is PHOSPHITE instead of phosphate. **PHOSCARE** is intended to be used as an immediate, highly available source of Phosphorous and Potassium.

Phosphorous is one of the most difficult nutrients to supply to plants, particularly through the soil because most soils have phosphorous fixation characteristics. To compensate for the inefficiency of phosphorous utilisation by plants it is necessary to apply **PHOSCARE** to the plants as it is an immediate form of Phosphorous which the plant recognises and utilises quickly.

PHOSCARE can be applied to ALL crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocados, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

GUARANTEED ANALYSIS

Total Product Basis	%w/w/v
Total Nitrogen (N)	0.0%
Total Phosphorous (P)	28.0%
Total Potassium (K)	25.0%

Phosphorous and Potassium present as mono acid dipotassium phosphite

KEEP OUT OF REACH OF CHILDREN

Always perform a compatibility test before mixing anything with **PHOSCARE**.

PHOSCARE is INCOMPATIBLE with trace elements, specifically Copper, Magnesium, and Calcium foliar nutrients.

CAUTION: Do not apply Copper for at least 30 days before or after a **PHOSCARE** application.

NET CONTENTS

3.785 L

Lot #

Directions and General Recommendations

SOIL APPLICATION

Apply **PHOSCARE** at 5.0 – 8.0 Litres per hectare through drip irrigation before flowering. Repeat as needed after flowering throughout the season.

FOLIAR APPLICATION

Apply **PHOSCARE** at 2.5 – 5.0 ml/ 1L of water and spray on to the foliage of the plants. Repeat as needed.

ALL FIELD CROPS (grain crops, flowers, etc...): Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** 2-3 weeks after crop emergence. Repeat the application every month.

ALL VEGETABLE CROPS: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** once every month starting 2-3 weeks after crop emergence.

POTATOES & CARROTS: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** six weeks after planting and once carrots are at 10-15 cm in height after emergence. Repeat at bulking stage. Repeat 2-3 weeks before harvest.

GREENHOUSE CROPS: Apply **PHOSCARE** at 3 ml/ 1L of water and spray on to the foliage of the plants. Repeat every month.

APPLES, POME & STONE FRUIT: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** before flowering, 1 week after fruit set. Repeat the application mid season. Repeat the application 2 weeks before harvest.

LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mistanding or any application not specifically described on the label.

FRUIT TREES & AVOCADOS: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** before flowering, 1 week after fruit set. Repeat the application mid season. Repeat the application 2 weeks before harvest.

KIWI FRUIT: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** before flowering, 1 week after fruit set. Repeat the application mid season. Repeat the application 2 weeks before harvest.

TABLEGRAPES & CITRUS: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** before flowering, 1 week after fruit set. Repeat the application mid season. Repeat the application 2 weeks before harvest.

WINEGRAPES: Apply 2.5 – 6.0 Litres per hectare of **PHOSCARE** before flowering, 1 week after fruit set. Repeat the application mid season. Repeat the application 2 weeks before harvest.

PHOSCARE is best applied when plants are in an active growing stage. **PHOSCARE** may be used at any time throughout the season. For best results and more specific recommendations, contact your nearest distributor.

Imported and Distributed in Australia exclusively by: Zaddo For Quality Gro PTY LTD, ABN 57080 750 084 | 9/9A Foundry Road, Seven Hills NSW 2147
Ph: 02 9838 9111 Fax: 02 9838 9110 | email: inquiries@zaddo.com.au website: www.zaddo.com.au Imported and Distributed in New Zealand exclusively by:
Roots Shoots and Fruits Ltd | PO Box 72, Waieke Island, Auckland, New Zealand | Ph: +64(0) 9372 9155 Fax: +64(0) 9372 9156 | email: rsf@rd2.co.nz
website: www.rd2.co.nz Manufactured by: JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA