

# BIOMIN<sup>TM</sup>

## BOOSTER V

Soluble Powder  
Multi-mineral Combination



### GUARANTEED ANALYSIS

	Dry Weight Basis %/w/w
<b>As amino acid Chelates:</b>	
Magnesium (Mg)	4.1
Zinc (Zn)	4.1
Iron (Fe)	2.6
Manganese (Mn)	2.6
Copper	1.5
Boron (B)	1.5
Molybdenum (Mo)	0.5
<b>As amino acid:</b>	
Nitrogen (N)	6.7

### GENERAL INFORMATION

**BioMin Booster V** is an amino acid chelated multi mineral. The chelating agent is mainly glycine, the smallest amino acid commonly used by and found in plants. The unique formulation of BioMin Booster V classifies it at the top of the range of all chelated minerals.

**BioMin Booster V** is a readily bio-available plant multi mineral product. The highlights of such a product include almost total absorption within a few hours after application. The chelating agent Glycine prevents the precipitation of the product and enables all the minerals to move freely inside the plant making the product highly systemic.

**BioMin Booster V** is ideal for all crops as it contains all the essential trace elements required for productivity, with additional Magnesium for chlorophyll production.

### ADVANTAGES & BENEFITS

**BioMin** products have the smallest molecular size that a chelate can ever have resulting in easy penetration of leaf surfaces.

**BioMin Booster V** aids in improving yields and crop quality when used according to the crops needs.

**BioMin Booster V** is very stable in formulation and can be used on all horticultural crops at almost any stage of growth

**BioMin Booster V** provides balanced nutrition supplying

all essential trace elements, including Molybdenum, a commonly overlooked essential trace element.

**BioMin Booster V** is 100% soluble.

**BioMin Booster V** is an excellent product for pre-flowering and post harvest application of elements to replenish lost reserves of trace elements and Magnesium in all crops, particularly tree and vine crops. This will enable improved bud fertility for bud burst and / or bud movement to proceed smoothly with all the necessary nutrients.

### COMPATIBILITY

ALWAYS run a compatibility test before spraying with other chemicals.

**BioMin Booster V** can also be used in combination with all other BioMin products, except BioMin Calcium.

**BioMin Booster V** is compatible with almost all fungicides, insecticides and herbicides.

**BioMin Booster V** is incompatible with Phosphorous, Calcium and Potassium foliar nutrients.

### PLANT & ENVIRONMENTAL SAFETY

**BioMin Booster V** is totally harmless to plants even when recommended rates are exceeded. Exceeding recommended rates is however unnecessary.

**BioMin Booster V** is totally harmless to both humans and wildlife and is environmentally friendly.

IMPORTED and DISTRIBUTED EXCLUSIVELY IN NEW ZEALAND BY

**ROOTS SHOOTS & FRUITS Ltd**

PO Box 72, Waiheke Island, New Zealand

Ph: +64(0)93729155 Fax: +64(0)93729156

E-mail: [rsf@rd2.co.nz](mailto:rsf@rd2.co.nz)

Web: <http://www.rd2.co.nz>




## APPLICATION GUIDELINES

CROP	TIME OF APPLICATION		RATE OF APPLICATION
APPLES & PEARS	1 <sup>st</sup> application 2 <sup>nd</sup> application	1-2 weeks before flowering 4-6 weeks before dormancy	1 – 1.5 kg/ha 1.5 – 2 kg/ha
AVOCADO	1 <sup>st</sup> application 2 <sup>nd</sup> application	1-2 weeks before flowering When fruit reaches walnut size	1 – 1.5 kg/ha 1 – 1.5 kg/ha
BRASSICAS and LETTUCE	1 <sup>st</sup> application 2 <sup>nd</sup> application	At 4-5 leaf stage 2-3 weeks later	1 – 1.5 kg/ha 1 – 1.5 kg/ha
GRAPES (TABLE, WINE, and DRIED FRUIT)	1 <sup>st</sup> application 2 <sup>nd</sup> application	1-2 weeks before flowering 2-3 weeks after harvest	1 – 1.5 kg/ha 1 – 1.5 kg/ha
KIWI GOLD(16A) & HAYWARD	1 <sup>st</sup> application 2 <sup>nd</sup> application	1-2 weeks before flowering 6-8 weeks before dormancy	1 – 1.5 kg/ha 1 – 1.5 kg/ha
POTATOES, ONIONS & CARROTS	1 <sup>st</sup> application 2 <sup>nd</sup> application	Once crop is 15 cm in height 1-2 weeks before bulking	1 – 1.5 kg/ha 1 – 1.5 kg/ha
STONE FRUIT & CHERRIES	1st application	3-4 weeks before leaf drop	1.5 – 2 kg/ha
STRAWBERRIES & ALL BERRIES	3 – 4 applications	Once every month beginning at 6 – 8 true leaf stage	1 – 1.5 kg/ha
TOMATO and CAPSICUM (& chillies)	3 – 4 applications	Once every month beginning once plants reach 15 – 20cm in height	1 – 1.5 kg/ha

For information on application rates and timing for crops not listed on this brochure, please contact your local distributor or visit the RSF website at [www.rd2.co.nz](http://www.rd2.co.nz)



It is always advisable that a leaf sample be taken before applying fertilisers to best ascertain the levels of elements and the nutrient requirements of the crop.



*Chelation is a natural process which occurs within plants. Elements are naturally chelated by plants into amino acid form before entry into the root system. These elements can only be utilised in this amino acid form in order to prevent absorbed nutrients from precipitation.*

*The same principle applies to foliar applied elements. Glycine chelates (amino acid) have been proven to be the most effective in supplying minerals to plants in order to correct nutrient deficiencies.*

*Glycine is the simplest amino acid with a molecular weight of 75. Chelates of glycine with cations have been fully studied and the picture to the left illustrates two moles of ligand (glycine) and one mole of metal forming a true chelate.*

*Above all, Glycine chelates have been proven to be the most effective, stable and economical products worldwide in the supply of plant nutrients.*

**IMPORTED and DISTRIBUTED EXCLUSIVELY IN AUSTRALIA BY ZADCO FOR QUALITY GRO P/L**  
 9/9A Foundry Road Seven Hills NSW 2147  
**Ph:** +61(02)98389111 **Fax:** +61(02)98389110  
 E-mail: [inquiries@zadco.com.au](mailto:inquiries@zadco.com.au)  
 Web: <http://www.zadco.com.au>

**MANUFACTURED BY**  
 JH Biotech, Inc.  
 4591 Olivas PK. Dr. Ventura  
 California 93006 USA  
 Web: <http://www.jhbiotech.com>