



FOLIAR NUTRITION PROGRAM

BRASSICAS



SOIL HEALTH · PLANT HEALTH · YOUR HEALTH

STAGE OF GROWTH	TASK	PRODUCT	VOLUME	NOTES
1st Application: (When plants reach 6-8 leaf stage)		Biomin Boron	1kg/ha	
		Biomin Booster V	1 kg /ha	
		Mobilizer	2L / ha	
2nd Application: (2 weeks later)		Biomin Calcium	1kg / Ha	
		Mobilizer	1L/ Ha	
3rd Application: (1 week later)		Biomin Boron	1.5 Kg / ha	
		Mobilizer	2L / Ha	
4TH Application: (around head formation)		Phoscare	2.5 L / Ha	
		Mobilizer	1L/ Ha	

SEEDLING DIP PREPARATION:

Superzyme: Soak 100-150 grams / 50 L of this beneficial inoculum in water overnight to activate it. Stir and Dip all seedlings, (including leaves) into the solution before planting. (Please ask for Superzyme protocols).

If the soil has been fumigated, or if there is a known problem in the soil, growers may consider beneficial inoculums be reapplied to the soil after fumigation traces have gone.

Inject through Irrigation: Superzyme 1kg/ha

Notes for all products in this program:

Use a wetter / sticking agent with every application.

Do not apply copper or copper containing products within 21 days before or after Phoscare application due to non compatibility.

This represents a guide only. It does not take into account certain microclimates, seasonal variations, or on farm management practises. Soil and leaf tests are always recommended to ascertain the nutritional status of crops to optimize product efficacy & plant performance.